

ANTIVENIN CROTALIDAE POLYVALENT EQUINE ORIGIN



RATTLERANTIVENIN™

Shown to be effective against envenomation due to North American Crotalidae.

Use: Rattler Antivenin is shown to be effective against envenomation in canine and equine due to North American Crotalidae. Crotalidae refers to cottonmouths/water moccasins, copperheads and rattlesnakes.

Dose: 50ml administered intravenously.

Administration: Keep frozen below -5°C (23°F) until use. Warm to body temperature using a warm water bath prior to administration. Administer to effect using a filtered IV set over a 20-60 minute period. Additional doses may be necessary. Do not reuse or refreeze after opening.

Administer Rattler Antivenin as is, without reconstitution or dilution. Additional fluids and supplemental therapies can be used, but should not be combined directly with Rattler Antivenin.

Age and Species: Canine: Safe for use eight weeks and older. Equine: Safe for use at any age.

Ingredients: Equine-derived Crotalidae Polyvalent Antivenin, Sodium Citrate

Caution: Every reasonable precaution has been taken to safeguard this product. While this serial has been reviewed and found to be safe, there is always the possibility, while rare, of adverse events. Please review your patient's history of receipt of equine protein products prior to administration.

See package insert for detailed product information.

Restricted to use by or under the direction of a licensed veterinarian.

serial:

expires:

Mg Biologics

U.S.Vet Lic. No. 614
(877) 769-2340 mgbiologics.com Ames, IA 50014